arXiv:math/0207296v1 [math.MG] 31 Jul 2002

Products of hyperbolic metric spaces

Thomas Foertsch* , Viktor Schroeder

Abstract

Let (X_i, d_i), $i = 1, 2$, be proper geodesic hyperbolic metric spaces. We give a general construction for a "hyperbolic product" $X_1 \times_h X_2$ which is itself a proper geodesic hyperbolic metric space and examine its boundary at infinity.

1 Introduction

Let (X_i, d_i), $i = 1, 2$, be proper geodesic hyperbolic metric spaces (for definitions see Section 2). We give a general construction for a "hyperbolic product" $X_1 \times_h X_2$ which is itself a proper geodesic hyperbolic metric space.
This construction only depends on chosen basepoints $z_i \in X_i$ or on points $u_i \in \partial X_i$, where ∂X_i is the boundary at infinity.

For given points $z_i \in X_i$ consider the set

$$Y \;:=\; \Big\{(x_1, x_2) \in X_1 \times X_2 \;\Big|\; d_1(x_1, z_1) \;=\; d_2(x_2, z_2)\Big\} \;\subset\; X_1 \times X_2$$

with the induced product metric

$$d_2\Big((x_1, x_2), (y_1, y_2)\Big) \;=\; \Big(d_1^2(x_1, y_1) \;+\; d_2^2(x_2, y_2)\Big)^{\frac{1}{2}}.$$

Let $d : Y \times Y \longrightarrow \mathbb{R}^+$ be the corresponding interior metric

$$d(x, y) \;:=\; \inf\Big\{L(c) \;\Big|\; c \text{ connecting } x \text{ to } y \text{ in } Y\Big\},$$

where $L(c)$ denotes the length of c in (Y, d_e).
We call (Y, d) the *hyperbolic product* of (X_1, z_1) and (X_2, z_2) and denote it also by

$$Y \;=\; (X_1, z_1) \times_h (X_2, z_2).$$

This notion is justified by the following

---

*supported by SNF Grant 21 - 589 38.99

2000 Mathematics Subject Classification. Primary 53C21

Theorem 1 *Let X_i, $i = 1, 2$, be proper geodesic hyperbolic spaces and $z_i \in X_i$, $i = 1, 2$. Then $Y = (X_1, z_1)\times_h(X_2, z_2)$ is also a proper geodesic hyperbolic space and ∂Y is naturally homeomorphic to $\partial X_1 \times \partial X_2$.*

The construction can be carried over in the limit case that the points z_i tend to infinity. This limit case seems to be of particular interest.
Let $u_i \in \partial X_i$ be given. These points give rise to Busemann functions $B_i : X_i \longrightarrow \mathbb{R}$. Define now

$$Y \ := \ \Big\{(x_1, x_2) \in X_1 \times X_2 \ \Big| \ B_1(x_1) \ = \ B_2(x_2)\Big\} \ \subset \ X_1 \times X_2$$

and consider as above the interior metric d on Y.
We call (Y, d) the hyperbolic product of (X_1, B_1) and (X_2, B_2), denote it by

$$Y \ = \ (X_1, B_1) \times_h (X_2, B_2),$$

and obtain the

Theorem 2 *Let X_i, $i = 1, 2$, be proper geodesic hyperbolic spaces and $B_i : X_i \longrightarrow \mathbb{R}$ Busemann functions on X_i. Then $Y = (X_1, B_1)\times_h(X_2, B_2)$ is also a proper geodesic hyperbolic space and ∂Y is naturally homeomorphic to the smashed product $\partial X_1 \wedge \partial X_2$.*

Remark 1 **i)** *The smashed product $\wedge$ is a standard construction for pointed topological spaces. Let (U_1, u_1), (U_2, u_2) be two pointed spaces then the smashed product $U_1 \wedge U_2$ is defined as $U_1 \times U_2/U_1 \vee U_2$, where $U_1 \times U_2$ is the usual product and*

$$U_1 \vee U_2 \ = \ \Big(\{u_1\} \times U_2\Big) \cup \Big(U_2 \times \{u_2\}\Big) \ \subset \ U_1 \times U_2$$

is the wedge product canonically embedded in $U_1 \times U_2$. Thus $U_1 \wedge U_2$ is obtained from $U_1 \times U_2$ by collapsing $U_1 \vee U_2$ to a point. For example $S^m \wedge S^n = S^{m+n}$.

ii) *In [BrFa] the authors proved that the hyperbolic product of real hyperbolic spaces $(\mathbb{H}^{m_1}, u_1)\times_h(\mathbb{H}^{m_2}, u_2)$, with u_i in the ideal boundary, is isometric to a real hyperbolic space $\mathbb{H}^{m_1+m_2-1}$.*
In [FS] the authors proved that, more generally, the hyperbolic product (Y, d) of Hadamard manifolds of pinched negative sectional curvature $-b^2 \leq K_i \leq -a^2 < 0$ is hyperbolic. This was done by showing that there exists a metric on Y that is bilipschitz to the one induced by the canonical embedding $i : Y \longrightarrow X$ and that carries pinched negative sectional curvature.
For related results also see [L1], [L2], [F1] and [F2].

iii) *Instead of the Euclidean product metric d_e on Y we could also take e.g. the maximum metric*

$$d_m\Big((x_1, x_2), (y_1, y_2)\Big) \ := \ \max\Big\{d_1(x_1, y_1), d_2(x_2, y_2)\Big\}$$

and the corresponding inner metric d' on Y. Note that d_e and d_m and therefore also d and d' are bilipschitz related.
The following holds in general: If (Y, d) is a proper geodesic metric space and d' is an other interior metric on Y which is bilipschitz related to d, then (Y, d') also is a proper metric space which implies that it is geodesic since (Y, d) is a length space (see e.g. Theorem 2.5.23 in [BuBuI]). In addition (Y, d) is hyperbolic if and only if (Y, d') is hyperbolic and in that case $\partial(Y, d)$ is homeomorphic to $\partial(Y, d')$.
For technical reasons we use in our proof the metric d_m on $X_1 \times X_2$.

iv) *Finally note that the hyperbolic product can similar be defined for finitely many factors and the analogue of Theorems 1 and 2 hold in that case.*

Outline of the paper:
In Section 2 we collect the necessary results on hyperbolic metric spaces. In Sections 3 and 4 we give a proof of Theorem 2. At the end of Section 3 we indicate the necessary changes for the situation of Theorem 1.

Acknowledgment: We want to thank Urs Lang for useful discussions.

2 Preliminaries

2.1 Hyperbolicity

A metric space (X, d) is called *geodesic*, if any two points $x, y \in X$ can be joined by a geodesic segment $\overline{xy}$ that is the image of a geodesic path $\gamma_{xy} : [0, d(x, y)] \longrightarrow X$ from x to y which is parameterized by arclength.
A geodesic metric space is called *δ-hyperbolic* if for any triangle with geodesic sides in X each side is contained in the δ- neighborhood of the union of the two other sides.
The space is called *hyperbolic* if it is δ-hyperbolic for some $\delta \geq 0$.

Let X be a metric space and $x, y, z \in X$. Then there exist unique $a, b, c \in \mathbb{R}_0^+$ such that

$$d(x, y) = a + b, \quad d(x, z) = a + c \quad \text{and} \quad d(y, z) = b + c.$$

In fact those numbers are given through

$$a \; = \; (y \cdot z)_x \;, \quad b \; = \; (x \cdot z)_y \;, \quad \text{and} \quad c \; = \; (x \cdot y)_z \;,$$

where for instance

$$(y \cdot z)_x \; = \; \frac{1}{2}\Big[d(y, x) \; + \; d(z, x) \; - \; d(y, z)\Big].$$

In the case that $X$is geodesic we may consider a geodesic triangle $\overline{xy} \cup \overline{xz} \cup \overline{yz} \subset X$, where for example $\overline{xy}$ denotes a geodesic segment connecting x to y. Given

such a triangle we denote by $\tilde{x} = \gamma_{xy}(a)$ the unique point on $\overline{yz}$ satisfying $d(\tilde{x}, y) = (x \cdot z)_y$ and in the same way we define $\tilde{y} \in \overline{xz}$ and $\tilde{z} \in \overline{xy}$.
Note that for X being a tree all these points coincide, i.e. $\tilde{x} = \tilde{y} = \tilde{z}$. In general an upper bound for the distances of these points measures the hyperbolicity of (X, d).

Lemma 1 **i)** *If (X, d) is δ-hyperbolic, then*

$$d(z, \tilde{z}) \ \le \ c + 2\delta, \qquad\qquad d\Big(\gamma_{xy}(t), \gamma_{xz}(t)\Big) \ \le \ 4\delta \quad \forall t \in [0, a]$$

and the points $\tilde{x}$, $\tilde{y}$, $\tilde{z}$ have pairwise distance $\le 4\delta$.

ii) *A metric space (X, d) is hyperbolic if and only if there exists a $\delta' \in \mathbb{R}_0^+$ such that given any geodesic triangle $\overline{xy} \cup \overline{xz} \cup \overline{yz} \subset X$ the points $\tilde{x} \in \overline{yz}$, $\tilde{y} \in \overline{xz}$ and $\tilde{z} \in \overline{xy}$ as defined above have distance less than δ' to each other.*

Proof: *i)* By δ-hyperbolicity $d(\tilde{z}, \overline{xz}) \ge \delta$ or $d(\tilde{z}, \overline{yz}) \ge \delta$. By triangle inequality we have in the first case $d(\tilde{z}, \tilde{y}) \le 2\delta$ and hence $d(\tilde{z}, z) \le c + 2\delta$. The other case is similar.
Assume that there is $t_0 \in [0, a]$ with $d(\gamma_{xy}(t_0), \gamma_{xz}(t_0)) > 4\delta$, then $d(\gamma_{xy}(t_0 - \delta), \gamma_{xz}(t_0 - \delta)) > 2\delta$ which implies $d(\gamma_{xy}(t_0 - \delta), \overline{xz}) > \delta$ and by hyperbolicity $d(\gamma_{xy}(t_0 - \delta), \overline{yz}) < \delta$. Let $p \in \overline{yz}$ be a point of minimal distance to $\gamma(t_0 - \delta)$. By triangle inequality $d(p, y) \ge b$ and hence $d(p, z) \le c$. Thus

$$\begin{aligned} a \ + \ c \ = \ d(x, z) \ &\le \ (t_0 - \delta) \ + \ d\Big(\gamma_{xy}(t_0 - \delta), p\Big) \ + \ d(p, z) \\ &< \ a \ + \ c \,; \end{aligned}$$

a contradiction. Since the corresponding estimate holds for the other sides as well the points $\tilde{x}$, $\tilde{y}$ and $\tilde{z}$ have pairwise distance $\le 4\delta$.
For *i)* compare to Proposition III.1.17 in [BriH]. □

2.2 T-functions

Definition 1 *Let $\alpha, \omega \in \mathbb{R}$ and $I := [\alpha, \omega]$.*

i) *A function $f : I \longrightarrow \mathbb{R}$ is called a T-function if f is continuous and there exists $\alpha + a \in [\alpha, \omega]$ such that the restrictions $f|_{(\alpha, \alpha+a)}$ and $f|_{(\alpha+a, \omega)}$ are differentiable with*

$$f'|_{(\alpha, \alpha+a)} \ \equiv \ -1 \qquad \textit{and} \qquad f'|_{(\alpha+a, \omega)} \ \equiv \ 1.$$

ii) *A function $f : I \longrightarrow \mathbb{R}$ is called a δ-T-function, $\delta \in \mathbb{R}_0^+$, if there exists a T-function $g : I \longrightarrow \mathbb{R}$ such that $||f - g||_{\sup} \ < \ \delta$.*

iii) *Let X be a geodesic metric space. A function $f : X \longrightarrow \mathbb{R}$ is called a δ-T-function, if for any geodesic segment $\gamma : [\alpha, \omega] \longrightarrow \mathbb{R}$ the function $f \circ \gamma$ is a δ-T-function.*

Remark 2 *It is straight forward to check that*

i) *Every T-function f is convex and Lipschitz with Lipschitz constant $l = 1$.*

ii) *For $[\alpha, \omega] \subset \mathbb{R}$ and $t_1, t_2 \in \mathbb{R}$ with $|t_1 - t_2| \leq |\alpha - \omega|$ there exists a unique T-function with $f(\alpha) = t_1$ and $f(\omega) = t_2$. Indeed there are unique $a, b, c \in \mathbb{R}_0^+$ such that $a + b = |\alpha - \omega|$, $a + c = t_1$, $b + c = t_2$. These are given via*

$$\begin{aligned} a &= \frac{1}{2}\Big(|\alpha - \omega| \;+\; t_1 \;-\; t_2\Big), \\ b &= \frac{1}{2}\Big(|\alpha - \omega| \;+\; t_2 \;-\; t_1\Big), \\ c &= \frac{1}{2}\Big(t_1 \;+\; t_2 \;-\; |\alpha - \omega|\Big). \end{aligned}$$

Now f satisfies

$$f(\alpha) \;=\; t_1, \qquad f(\alpha + a) \;=\; c, \qquad f(\omega) \;=\; t_2.$$

iii) *f is a (δ-)T-function $\Longrightarrow$ $f + const$ is a (δ-)T-function.*

iv) *A limit of a sequence of (δ-)T-functions is a (δ-)T-function.*

Lemma 2 *Let X be a geodesic metric space. Then the following are equivalent:*

i) *X is hyperbolic.*

ii) *There exists a $\delta \in \mathbb{R}^+$ such that for all $x \in X$ the function*

$$d_x : X \longrightarrow \mathbb{R}\,, \qquad y \longmapsto d(x, y)$$

is a δ-T-function.

Proof: "$\Longrightarrow$" Let X be δ-hyperbolic, $x, y, z \in X$, $d(x, z) = a + c$, $d(y, z) = b + c$ and $\gamma : [0, a + b] \longrightarrow \overline{xy}$ an arc length parameterized geodesic connecting x to y.
Now consider the T-function $f : [0, a + b] \longrightarrow \mathbb{R}^+$ determined by

$$f(0) = d(z, \gamma(0)) = a + c \quad \text{ and } \quad f(a + b) = d(z, \gamma(a + b)) = b + c.$$

Note that $f(a) = c$ and $(d_z \circ \gamma)(a) \leq c + 4\delta$ by Lemma 1 *ii*).
From the fact that $d_z \circ \gamma : [0, a + b] \longrightarrow \mathbb{R}^+$ is 1-Lipschitz it immediately follows that

$$||d_z \circ \gamma \;-\; f||_{\sup} \;\leq\; 4\delta.$$

Hence $d_z \circ \gamma$ is a 4δ-T-function.
"$\Longleftarrow$" Let now X satisfy condition *ii*). We show that X is hyperbolic using the criterion of Lemma 1 *i*):

For $x, y, z \in X$ and geodesic segments $\overline{xy}$, $\overline{xz}$ and $\overline{yz}$ connecting these points, condition *ii)* yields $c \leq d(z, \tilde{z}) \leq c + \delta$. We now consider the geodesic triangle $\overline{x\tilde{z}} \cup \overline{z\tilde{z}} \cup \overline{xz}$, where $\overline{x\tilde{z}} \subset \overline{xy}$ and $\overline{z\tilde{z}}$ is any geodesic segment connecting z to $\tilde{z}$. For $\hat{z} \in \overline{xz}$ satisfying $d(z, \hat{z}) = (x \cdot \tilde{z})_z$ condition *ii)* gives $d(\tilde{z}, \hat{z}) < \delta$. Furthermore one has $c < d(z, \hat{z}) < c + \frac{\delta}{2}$ and therefore $d(\tilde{y}, \hat{z}) < \frac{\delta}{2}$. Thus we achieve

$$d(\tilde{z}, \tilde{y}) \;\leq\; d(\tilde{z}, \hat{z}) \;+\; d(\hat{z}, \tilde{y}) \;\leq\; \delta \;+\; \frac{\delta}{2} \;=\; \frac{3}{2}\delta.$$

The same argument of course yields $d(\tilde{z}, \tilde{x}) \leq \frac{3}{2}\delta$ and $d(\tilde{y}, \tilde{x}) \leq \frac{3}{2}\delta$. By Lemma 1 *ii)* we obtain the result. $\square$



Figure 1: The graph of $d_z \circ \gamma$ lies in the region between the graphs of f and $f + \delta$.

2.3 The boundary at infinity and Busemann functions

We associate to a hyperbolic space (X, d) a boundary ∂X at infinity. There are different descriptions in the literature (see e.g. [BeKa]) all of which coincide for proper geodesic hyperbolic spaces.

We choose a basepoint $z \in X$. We say that a sequence $\{x^k\}_{k \in \mathbb{N}}$ of points in X *converges to infinity*, if

$$\liminf_{k,l \longrightarrow \infty} \; (x^k \cdot x^l)_z \;=\; \infty.$$

Two sequences $\{x^k\}_{k \in \mathbb{N}}$ and $\{y^k\}_{k \in \mathbb{N}}$ converging to infinity are equivalent, $\{x^k\}_{k \in \mathbb{N}} \sim \{y^k\}_{k \in \mathbb{N}}$ if

$$\liminf_{k,l \longrightarrow \infty} \; (x^k \cdot y^l)_z \;=\; \infty.$$

One shows that $\sim$ is an equivalence relation and defines ∂X as the set of equivalence classes. We write $[\{x^k\}] \in \partial X$ for the corresponding class. One can also show that for every $x \in X$ and $v \in \partial X$ there is a geodesic ray $\gamma_{xv} : [0, \infty] \longrightarrow X$ parameterized by arclength with $\gamma_{xv}(0) = x$ and $[\{\gamma_{xv}(k)\}] = v$.

For $v \in \partial X$ and $r > 0$ one defines

$$U(v, r) \ := \ \Big\{ w \in \partial X \ \Big| \ \exists \{x^k\}, \{y^k\} \text{ s.t. } [\{x^k\}] = v, \ [\{y^k\}] = w, \ \liminf_{k,l \longrightarrow \infty} \ (x^k \cdot y^l)_z \ \geq \ r \Big\}.$$

On ∂X we consider the topology generated by $U(v, r)$, $v \in \partial X$, $r > 0$.

If X is proper, ∂X is a compact topological space. We now also fix a basepoint $u \in \partial X$ and a geodesic ray γ_{zu} from z to u. The function $B : X \longrightarrow \mathbb{R}$, $B(x) := \lim_{t \to \infty} [d(x, \gamma_{zu}(t)) - t]$ is called the *Busemann function associated to* γ_{zu}. B is a δ-T-function as a limit of δ-T-functions.

Definition 2 *A geodesic ray* $\gamma : [0, \infty) \longrightarrow X$ *is called a* B*-ray if and only if* γ *is parameterized by arc length and*

$$B\Big(\gamma(t)\Big) \ = \ B\Big(\gamma(0)\Big) \ - \ t \qquad \forall t \in [0, \infty).$$

By a standard limit argument we obtain the

Lemma 3 *Let* X *be a complete, locally compact, hyperbolic metric space and* B *a Busemann function on* X*. Then for every* $x \in X$ *there exists a* B*-ray* γ *with* $\gamma(0) = x$*.*

Let now $v \in \partial X \setminus \{u\}$ and consider B-rays $\gamma_{\gamma_{zv}(t)}$ starting at $\gamma_{zv}(t)$. Then these rays subconverge as sets to a geodesic from v to u and by suitable reparameterization we obtain the existence of a geodesic $\gamma_{vu} : \mathbb{R} \longrightarrow X$ with

$$B\Big(\gamma_{vu}(t)\Big) \ = \ -t, \qquad \Big[\{\gamma_{vu}(k)\}\Big] \ = \ u \quad \text{ and } \quad \Big[\{\gamma_{vu}(-k)\}\Big] \ = \ v.$$

The hyperbolicity of X also implies that there exists a constant C (depending on u, v, z) such that

$$d\Big(\gamma_z(t), \gamma_{vu}(t)\Big) \ \leq \ C \quad \text{ and } \quad d\Big(\gamma_{zv}(t), \gamma_{vu}(-t)\Big) \ \leq \ C \qquad \forall t \geq 0.$$

Let now $x, y \in X$ and γ_x, γ_y be B-rays starting at x, y. Then γ_x, γ_y and a geodesic segment $\overline{xy}$ form an ideal triangle with vertices x, y and $u = \gamma_{zu}(\infty) \in \partial X$. We want to look for points $\tilde{u}, \tilde{x}, \tilde{y}$ as for finite triangles. Clearly there are $a, b \geq 0$ such that $d(x, y) = a + b$ and $B(\gamma_x(a)) = B(\gamma_x(b))$. Indeed

$$\begin{aligned} a \ &= \ (y \cdot B)_x \ \ := \ \ \frac{1}{2}\Big(d(y, x) \ + \ B(x) \ - \ B(y)\Big) \quad \text{and} \\ b \ &= \ (x \cdot B)_y \ \ := \ \ \frac{1}{2}\Big(d(x, y) \ + \ B(y) \ - \ B(x)\Big). \end{aligned}$$

Let $\tilde{u} \in \overline{xy}$ be the point with $d(x, \tilde{u}) = a$ and let $\tilde{y} = \gamma_x(a)$, $\tilde{x} = \gamma_y(b)$.



Figure 2: Triangle and ideal triangle in the hyperbolic space.

Lemma 4 **i)** *The points $\tilde{x}$, $\tilde{y}$ and $\tilde{u}$ have pairwise distance $\le 8\delta$.*

ii) *For all $t \ge 0$ it holds $d(\gamma_x(a+t), \gamma_y(b+t)) \le 8\delta$.*

Proof: Let $B(x) = \lim_{t\to\infty}(d(x, \gamma_{zu}(t)) - t)$, $z_i = \gamma_{zu}(i)$, $i \in \mathbb{N}$, and consider the triangle x, y, z_i with corresponding values $a_i, b_i, c_i \in \mathbb{R}^+$ as well as points

$$\hat{z}_i \;=\; \gamma_{xy}(a_i), \qquad \hat{y}_i \;=\; \gamma_{xz_i}(a_i) \qquad \text{and} \qquad \hat{x}_i \;=\; \gamma_{yz_i}(b_i).$$

Clearly one has $a_i \longrightarrow a$ and $b_i \longrightarrow b$.
Consider also the triangles $x, \gamma_x(a_i), z_i$. Note that $|d(z_i, \gamma_x(a_i)) - d(z_i, \hat{y}_i)| \longrightarrow 0$ which implies by Lemma 1 *ii)*

$$\limsup_{i\longrightarrow\infty} d\Big(\gamma_x(a_i), \tilde{y}_i\Big) \;\le\; 2\delta.$$

In the same way we obtain

$$\limsup_{i\longrightarrow\infty} d\Big(\gamma_y(b_i), \tilde{x}_i\Big) \;\le\; 2\delta.$$

Since $d(\tilde{x}_i, \tilde{y}_i) \le 4\delta$ by Lemma 1 we obtain *i)*.
The proof of *ii)* is similar. □

We need the following

Lemma 5 *Let $x, y \in X$ and $\sigma : [0, d(x,y)] \longrightarrow X$ a curve parameterized by arclength such that $\sigma(0) = x$ and $d(\sigma(d(x,y)), y) \le R$. Then*

$$d\Big(\gamma_{xy}(t), \sigma(t)\Big) \;\le\; \frac{3}{2}R \;+\; 4\delta \quad \forall t \in [0, d(x,y)].$$

Proof: For fixed $t \in [0, d(x,y)]$ consider the triangle $x, y, z := \sigma(t)$. Since $d(x,z) \le t$ and $d(z,y) \le d(x,y) - t + R$ we have $c = (x \cdot y)_z \le \frac{R}{2}$ and hence there exists $\tilde{z} = \gamma_{xy}(t')$ with $d(z, \tilde{z}) \le \frac{R}{2} + 4\delta$. Note that $t' = d(x,z) - c$ and $d(x,y) - t' = d(z,y) - c$. Thus $|t - t'| \le R$ from which the claim follows. □

2.4 A Morse estimate

We need an estimate whose proof is similar to the proof of the Morse inequality.

Lemma 6 *Let* (X,d) *be* δ*-hyperbolic,* $x,y \in X$ *and* $\gamma : [0,1] \longrightarrow X$ *be a continuous path from* x *to* y*. If there exists a point* $p = \gamma_{xy}(s_0) \in \overline{xy}$ *such that* $d(p,\gamma(t)) \geq R$ *for all* $t \in [0,1]$ *and* $R > 90\delta$*, then*

$$L(\gamma) \;\geq\; d(x,y) \;+\; \frac{1}{20\delta}R^2.$$

Proof: Define $a(t) := (y \cdot \gamma(t))_x \in [0, d(x,y)]$. Since a is continuous, $a(0) = 0$ and $a(1) = d(x,y)$ there are $0 < t_- < t_+ < 1$ such that

$$a(t_-) \;=\; s_0 \;-\; \frac{R}{2}\,, \qquad a(t_+) \;=\; s_0 \;+\; \frac{R}{2}\,, \qquad a(t) \in \Big[s_0 - \frac{R}{2}, s_0 + \frac{R}{2}\Big] \quad \forall t_- < t < t_+.$$

Choose $k+1 := [\frac{R}{8\delta}] + 1$ points $s_1 \leq ... \leq s_{k+1} \in [s_0 - \frac{R}{2}, s_0 + \frac{R}{2}]$ such that $|s_{i+1} - s_i| \geq 12\delta$ and let $t_i \in [t_-, t_+]$ be points $t_1 \leq t_2 \leq ... \leq t_{k+1}$ with $a(t_i) = s_i$. Now

$$L(\gamma) \;\geq\; d\Big(x, \gamma(t_-)\Big) \;+\; \sum_{i=1}^{k} d\Big(\gamma(t_{i+1}), \gamma(t_i)\Big) \;+\; d\Big(\gamma(t_+), y\Big).$$

By construction

$$d\Big(x, \gamma(t_-)\Big) \geq a(t_-) \qquad \text{and} \qquad d\Big(y, \gamma(t_+)\Big) \geq d(x,y) \;-\; a(t_-).$$

Thus $d(x,\gamma(t_-)) + d(\gamma(t_+), y) \geq d(x,y) - R$.
Let $q_i \in \overline{xy}$ be a point such that $d(\gamma(t_i), q_i) = d(\gamma(t_i), \overline{xy})$. Since $d_{\gamma(t_i)} \circ \gamma_{xy}$ is a 4δ-T-function by Lemma 2, a minimum of this function is assumed in distance $\leq 4\delta$ of the corresponding T-function. Thus $d(q_i, \gamma_{xy}(s_i)) \leq 4\delta$, which implies $d(q_i, q_{i+1}) \geq 4\delta$ since $d(\gamma_{x,y}(s_{i+1}), \gamma_{xy}(s_i)) \geq 12\delta$.
By Lemma 8.4.23 in [BuBuI] we obtain

$$d\Big(\gamma(t_{i+1}, \gamma(t_i))\;\geq\; d\Big(\gamma(t_{i+1}), q_{i+1}\Big) \;+\; d\Big(\gamma(t_i), q_i\Big) \;-\; 4\delta.$$

Since $d(\gamma(t_i), p) \geq R$ and $d(q_i, p) \leq \frac{R}{2} + 4\delta$ we obtain $d(\gamma(t_{i+1}), \gamma(t_i)) \geq R - 12\delta$. Thus

$$\begin{aligned} L(\gamma) &\geq d(x,y) \;+\; \Big[\frac{R}{12\delta}\Big](R \;-\; 12\delta) \;-\; R \\ &\geq d(x,y) \;+\; \frac{1}{20\delta}R^2 \end{aligned}$$

for $R \geq 90\delta$. □



Figure 3: This figure shows how Lemma 8.4.23 in [BuBuI] is applied to the segments $\gamma_{xy}|_{[s_i,s_{i+1}]}$ and $\gamma|_{t_i,t_{i+1}}$ in the proof of Lemma 6.

3 The hyperbolic product

In this section we prove the first part of Theorem 2, which is equal to Proposition 3.

Let (X_i, d_i) be δ_i-hyperbolic spaces, $i = 1, 2$, and $\delta := \max\{\delta_1, \delta_2\}$. Let further $B_i : X_i \longrightarrow \mathbb{R}$ be Busemann functions on X_i. We study the set

$$Y \;:=\; \Big\{(x_1, x_2) \in X_1 \times X_2 \;\Big|\; B_1(x_1) \;=\; B_2(x_2)\Big\}.$$

On Y we consider the maximum metric $d_m : Y \times Y \longrightarrow \mathbb{R}$,

$$d_m\Big((x_1, x_2), (x_1', x_2')\Big) \;:=\; \max\Big\{d_1(x_1, x_1'), d_2(x_2, x_2')\Big\}.$$

For our use of d_m instead of d_e compare Remark 1 *iii*) in the Introduction.
Let $p, p' \in Y$. We first construct two curves in Y between p and p', the Γ-*curve* $\Gamma_{pp'}$ and the *continuous* Γ-*curve* $\Gamma^c_{pp'}$.
The advantage of $\Gamma_{pp'}$ is that this curve is conceptually easy to understand. However, $\Gamma_{pp'}$ is in general not continuous, $\Gamma^c_{pp'}$ is a continuous variation of $\Gamma_{pp'}$.
Let $p = (p_1, p_2)$, $p' = (p_1', p_2')$ and γ_i, γ_i' be B_i-rays starting at p_i, p_i'. Let further $\gamma, \gamma' : \mathbb{R}^+_0 \longrightarrow Y$ be the geodesic rays

$$\gamma(t) \;=\; \Big(\gamma_1(t), \gamma_2(t)\Big)\;, \qquad \gamma'(t) \;=\; \Big(\gamma_1'(t), \gamma_2'(t)\Big).$$

These geodesics are parameterized by constant speed 1. We set

$$\begin{aligned} a_i \;:=\; (p_i' \cdot B_i)_{p_i} \;&=\; \frac{1}{2}\Big(d(p_i, p_i') \;+\; B_i(p_i) \;-\; B_i(p_i')\Big) \quad \text{and} \\ b_i \;:=\; (p_i \cdot B_i)_{p_i'} \;&=\; \frac{1}{2}\Big(d(p_i, p_i') \;+\; B_i(p_i') \;-\; B_i(p_i)\Big)\;, \end{aligned}$$

such that $a_i + b_i = d_i(p_i, p'_i)$. Let $a := \max\{a_1, a_2\}$ and $b := \max\{b_1, b_2\}$, then $a + b = d_m(p, p')$. We define by a slight abuse of notation

$$\Gamma \; = \; \Gamma_{pp'} \; := \; \gamma|_{[0,a]} \; * \; {\gamma'|_{[0,b]}}^{-1}.$$

Note that $\Gamma_{pp'}$ is not necessarily continuous, since $\gamma(a) = (\gamma_1(a), \gamma_2(a))$ is not necessarily equal to $\gamma'(b) = (\gamma'_1(b), \gamma'_2(b))$. However, $d_i(\gamma_i(a), \gamma'_i(b)) \le 8\delta_i$ by Lemma 4.

The curve $\Gamma^c_{pp'}$ is a continuous modification of $\Gamma_{pp'}$ and defined as

$$\Gamma^c_{pp'} \; := \; \gamma|_{[0,a+2\delta]} \; * \; \Gamma_1 \; * \; \Gamma_2 \; * \; {\gamma'|_{[0,b+2\delta]}}^{-1},$$

where Γ_1 is a continuous curve in Y form $(\gamma_1(a+2\delta), \gamma_2(a+2\delta))$ to $(\gamma'_1(b+2\delta), \gamma_2(a+2\delta))$ and Γ_2 a continuous curve in Y from $(\gamma'_1(b+2\delta), \gamma_2(a+2\delta))$ to $(\gamma'_1(b+2\delta), \gamma'_2(b+2\delta))$ given in the following way:
Let $\eta_1 : [\alpha_1, \beta_1] \longrightarrow X_1$ be a geodesic from $\gamma_1(a+2\delta)$ to $\gamma'_1(b+2\delta)$. Note that $L(\eta_1) \le 8\delta$ and $B_1(\eta_1(\alpha_1)) = B_2(\eta_1(\alpha_2))$. Since B_1 is 1-Lipschitz and a 4δ-T-function we obtain

$$B_1\Big(\eta_1(t)\Big) \; \le \; B_1\Big(\eta_1(\alpha_1)\Big) \; + \; 2\delta \; = \; B_1(p_1) \; - \; a.$$

Thus $-B_1(\eta_1(t)) + B_1(p_1) \ge a \ge 0$ and

$$\Gamma_1(t) \; = \; \Big(\eta_1(t), \gamma_2\Big(- B_1(\eta_1(t)) + B_1(p_1)\Big)\Big)$$

is well defined. By construction $\Gamma_1(t) \in Y$ and $L_{d_m}(\Gamma_1) \le 8\delta$.
In a similar way one constructs Γ_2.

We can easily estimate the length of $\Gamma^c_{pp'}$ and obtain the

Lemma 7 *Given two points $p, p' \in Y$ the continuous curve $\Gamma^c = \Gamma^c_{pp'}$ has length*

$$L(\Gamma^c) \; \le \; d_m(p, p') \; + \; 20\delta.$$

This immediately implies the

Proposition 1 *Given two points $p, p' \in Y$, it holds*

$$d_m(p, p') \; \le \; d(p, p') \; \le \; d_m(p, p') \; + \; 20\delta.$$

Thus (Y, d_m) and (Y, d) are quasi-isometric and hence bilipschitz on a large scale. But also on a local scale they induce the same topology:

Lemma 8 *The metrics d and $d_m|_Y$ induce the same topology on Y.*

Proof: We need to show that

$$\lim_{i \longrightarrow \infty} d_m|_Y(y_i, y_0) \; = \; 0 \qquad \Longrightarrow \qquad \lim_{i \longrightarrow \infty} d(y_i, y_0) \; = \; 0.$$

Thus it suffices to prove the

Sublemma 1 *Let $\{y_i\}_{i\in\mathbb{N}}$ be a sequence in Y that converges to $y_0 \in Y$ with respect to $d_e|_Y$. Then for all $\epsilon > 0$ there exists $\rho(\epsilon) > 0$ such that for all $y_i \in Y$ satisfying $d_e|_Y(y_i, y_0) < \rho$ there exists a curve Γ_i in Y connecting y_i to y_0 of length $L(\Gamma_i) \leq \epsilon$.*

A sequence $\{y_i\}_{i\in\mathbb{N}} = \{(y_{i1}, y_{i2})\}_{i\in\mathbb{N}}$ in $Y \subset X_1 \times X_2$ converges with respect to $d_m|_Y$ if and only if the sequences $\{y_{ij}\}_{i\in\mathbb{N}}$ in X_j converge with respect to d_j, $j = 1, 2$.
Define

$$K_j \ := \ \Big\{\gamma(\frac{\epsilon}{4}) \ \Big| \ \gamma \text{ is a } B_j\text{-ray with } \gamma(0) \ = \ y_{0j}\Big\}$$

and let γ_{ij} be B_j-rays with $\gamma_{ij}(0) = y_{ij}$. The local compactness of X_j implies the
Claim: There exists $\rho > 0$ such that $d_j(y_{ij}, y_{0j}) < \delta$ implies $d(\gamma_{ij}(\frac{\epsilon}{4}), K_j) \leq \frac{\epsilon}{8}$.

Let now $d_m(y_i, y_0) < \rho$, then $d_j(y_{ij}, y_{0j}) < \rho$ and thus by the claim there are B_j-rays γ_j starting at y_{0j} such that $d(\gamma_{ij}(\frac{\epsilon}{4}), \gamma_j(\frac{\epsilon}{4})) < \frac{\epsilon}{8}$. Similar to the construction of $\Gamma^c_{pp'}$ one now finds a continuous path $\tilde{\Gamma}^c$ in Y connecting successively the points

$$\Big(y_{i1}, y_{i2}\Big), \ \ \Big(\gamma_{i1}(\frac{\epsilon}{8}), \gamma_{i2}(\frac{\epsilon}{8})\Big), \ \ \Big(\gamma_1(\frac{\epsilon}{8}), \gamma_{i2}(\frac{\epsilon}{8})\Big), \ \ \Big(\gamma_1(\frac{\epsilon}{8}), \gamma_2(\frac{\epsilon}{8})\Big), \ \ \Big(y_{01}, y_{02}\Big)$$

of length

$$\tilde{\Gamma}^c \ < \ \frac{\epsilon}{4} \ + \ \frac{\epsilon}{8} \ + \ \frac{\epsilon}{8} \ + \ \frac{\epsilon}{4} \ = \epsilon.$$

□

Corollary 1 *(Y, d) is locally compact, complete and hence proper and geodesic.*

Proof: Since the Busemann functions b_i, $i = 1, 2$, are continuous, Y is a closed subset of the locally compact space $(X_1 \times X_2, d_m)$ and therefore locally compact itself when endowed with the induced metric $d_m|_Y$. Thus from Lemma 8 it follows that (Y, d) is also locally compact.
Every Cauchy-sequence in (Y, d) is a Cauchy-sequence in $(Y, d_m|_Y)$. But $(Y, d_m|_Y)$ is complete and therefore the Cauchy-sequence converges in $(Y, d_m|_Y)$. Now the proof of Lemma 8 yields convergence in (Y, d). Hence (Y, d) is complete.
Finally every locally compact, complete length space is proper and geodesic (see e.g. Proposition I.3.7. in [BriH]). □

Let $p, p' \in Y$ and let

$$\sigma : [0, d(p, p')] \longrightarrow Y \ , \qquad \sigma(t) \ = \ \Big(\sigma_1(t), \sigma_2(t)\Big)$$

be a unit speed geodesic from p to p'. We want to compare σ with the curve $\Gamma(p, p')$. To have the same domain, we modify $\Gamma_{pp'}$ a little: Let a, b as above and

$$\begin{aligned} a^* &= a + \frac{1}{2}\Big(d(p,p') - d_m(p,p')\Big) , \\ b^* &= b + \frac{1}{2}\Big(d(p,p') - d_m(p,p')\Big). \end{aligned}$$

We define $\Gamma^* = \Gamma^*_{pp'} : [0, d(p,p')] \longrightarrow Y$ via

$$\Gamma^*_{pp'}(t) := \gamma|_{[0,a^*]} * \gamma'|_{[0,b^*]}{}^{-1}$$

and prove the

Proposition 2 *For σ and Γ^* as above it holds*

$$d\Big(\sigma(t), \Gamma^*(t)\Big) \le 500\delta.$$

To simplify the arguments, we will assume:

(a) $d(p_1,p_1') \ge d(p_2,p_2')$ which implies $a = a_1$ and $b = b_1$,

(b) $B_i(p_i) = 0$, $B_i(\gamma_i(t)) = -t$, and $B_i(\gamma_i'(t)) = -t + (b-a)$.

We can assume this without loss of generality: (a) by interchanging the factors and (b) by adding the same constant to both Busemann functions.

The first step in the proof of Proposition 2 is the

Lemma 9 *There exists $t_0 \in [0, d(p,p')]$ with $d_1(\sigma_1(t_0), \gamma_1(a)) \le 30\delta$.*

Proof: Consider the ideal triangle $p_1, p_1', u_1 = \gamma_1(\infty) = \gamma_1'(\infty)$ in X_1 with points $\tilde p_1' = \gamma_1(a)$, $\tilde p_1 = \gamma'(b)$, $\tilde u_1 = \gamma_{p_1,p_1'}(a) \in \overline{p_1,p_1'}$ of pairwise distance $\le 8\delta$. Choose t_0 such that

$$M := d\Big(\sigma_1(t_0), \tilde u_1\Big) = \min_t d\Big(\sigma_1(t), \tilde u_1(t)\Big).$$

By Lemma 6 we have $L_{d_1}(\sigma_1) \ge (a+b) + \frac{1}{20\delta}M^2$. Since $L_{d_1}(\sigma_1) \le L_{d_m}(\sigma) \le (a+b) + 20\delta$ we obtain $M \le 20\delta$ and thus the result. □

We decompose σ into two pieces $\sigma = \bar\alpha * \bar\beta$ where $\bar\alpha = \sigma|_{[0,t_0]}$ and $\bar\beta = \sigma|_{[t_0, d(p,p')]}$, write $\bar\alpha = (\bar\alpha_1, \bar\alpha_2)$ and $\bar\beta = (\bar\beta_1, \bar\beta_2)$ and prove the

Lemma 10 *With the notation above it holds*

$$\Big|L(\bar\alpha_i) - a\Big| \le 50\delta \quad \text{and} \quad \Big|L(\bar\beta_i) - b\Big| \le 50\delta \quad \text{for } i = 1,2.$$

Proof: With M as in Lemma 9 we compute

$$L(\bar{\alpha}_i) \;\geq\; \Big|B_i\Big(\sigma_i(t_0)\Big) \;-\; B_i(p_i)\Big| \;=\; \Big|B_1\Big(\sigma_1(t_0)\Big)\Big| \;\geq\; a \;-\; M \;-8\delta \;\geq\; a \;-\; 30\delta.$$

and

$$L(\bar{\beta}_i) \;\geq\; \Big|B_i\Big(\sigma_i(t_0)\Big) \;-\; B_i(p_i')\Big| \;=\; \Big|B_1\Big(\sigma_1(t_0)\Big) \;-\; (b-a)\Big| \;\geq\; b \;-\; 30\delta.$$

Since $L(\bar{\alpha}_i) + L(\bar{\beta}_i) \leq a + b + 20\delta$ by Lemma 7 we obtain the result. □

Lemma 11 *With t_0 as in Lemma 9 it holds $d_2(\sigma_2(t_0), \gamma_2(a)) \leq 100\delta$.*

Proof: Consider the ideal triangle $p_2, q = \sigma_2(t_0) = \bar{\alpha}_2(t_0), u_2 = \gamma_2(\infty)$ in X_2 with corresponding points $\tilde{q} = \gamma_2((q \cdot B_2)_{p_2}), \tilde{u}_2, \tilde{p}_2$. Since

$$(q \cdot B_2)_{p_2} \;=\; \frac{1}{2}\Big(d(p_2, \bar{\alpha}_2(t_0)) \;+\; B_2(p_2) \;-\; B_2(\bar{\alpha}_2(t_0))\Big)$$

and $|L(\bar{\alpha}_1 - a)| \leq 50\delta$, $|B_2(p_2) - B_2(\bar{\alpha}_e(t_0))| \geq a - 30\delta$ by Lemma 10, we see

$$\Big|(q \cdot B_2)_{p_2} \;-\; a\Big| \;\leq\; 40\delta$$

and

$$d(\tilde{u}_2, \tilde{p}_2) \;=\; \frac{1}{2}\Big(d(p_2, \bar{\alpha}_2(t_0)) \;+\; B_2(\bar{\alpha}_2(t_0)) \;-\; B_2(p_2)\Big) \;\leq\; 40\delta.$$

Together with Lemma 4 we get the estimate. □

Proof of Proposition 2: Lemmata 9, 11 and Proposition 1 imply that $d(\sigma(t_0), \gamma(a)) \leq 120\delta$. Combining some triangle inequalities we obtain $|t_0 - a| \leq 150\delta$ and

$$d\Big(\sigma(a^*), \gamma(a^*)\Big) \;\leq 300\delta\;, \qquad d\Big(\sigma(a^*), \gamma(b^*)\Big) \;\leq 300\delta.$$

Together with Lemma 5 we obtain Proposition 2. □

Proposition 3 *(Y, d) is hyperbolic.*

Proof: By Lemma 2 it suffices to show that there exists a Δ such that for all $q, p, p' \in Y$ and all minimal geodesics $\sigma : [0, d(p, p')] \longrightarrow Y$ from p to p' the function $t \longmapsto d(q, \sigma(t))$ is a Δ-T-function. By Propositions 1 and 2 it suffices to show that there exists Δ such that for all q, p, p' the function

$$[a^*, b^*] \longrightarrow \mathbb{R} \qquad t \longmapsto d_m\Big(q, \Gamma^*(t)\Big)$$

is a Δ-T-function, where $\Gamma^* = \gamma|_{[0,a*]} * \gamma'|_{[0,b^*]}{}^{-1}$ as defined above.
Define $f, g : [a^*, b^*] \longrightarrow \mathbb{R}$ via

$$f(t) \;:=\; d_1\Big(q_1, \Gamma_1^*(t)\Big) \quad \text{and} \quad g(t) \;:=\; d_2\Big(q_2, \Gamma_2^*(t)\Big).$$

We have to show that $\max\{f,g\}$ is a Δ-T-function. We use without loss of generality as above that $d_1(p_1,p_1') \geq d_2(p_2,p_2')$, i.e. that $d_m(p,p') = d_1(p_1,p_1')$. In this case $d(\Gamma_1(t),\gamma_{p,p'}(t)) \leq \delta'$ for all $t \in [0,a+b]$, where δ' only depends on δ and not on p,p'. It follows that f is a Δ-T-function for some Δ only depending on δ.

Note further that $g|_{[0,a^*]}$ and $g|_{[a^*+\epsilon,a^*+b^*]}$ are 4δ-T-functions by Lemma 2 for every $\epsilon > 0$, and hence $g|_{[a^*,a^*+b^*]}$ is a 12δ-T-function since the jump at a^* is bounded by 8δ.

Let $v = (q_2 \cdot B_2)_{p_2}$, then the function

$$[0,\infty) \longrightarrow [0,\infty)\ , \qquad t \longmapsto d_2\Big(q_2,\gamma_2(t)\Big)$$

assumes the minimum 8δ-close to the point $t = v$.

If $v > a$, hence $v \geq a^* - 10\delta$, then by the properties of δ-T-functions $g|_{[0,a^*+b^*]}$ is easily checked to be a 30δ-T-function.

Let us assume that $v < a$. Set $u := d_2(q_2,p_2) - v$ and let $\tilde{\gamma}_2 : [0,\infty) \longrightarrow X$ be a B_2-ray starting at q_2. Then $d_2(\tilde{\gamma}_2(u+t),\gamma_2(v+t)) \leq 8\delta$. In particular $d_2(\tilde{\gamma}_2(u+(a^*-v)),\gamma_2(a^*)) \leq 8\delta$ which implies

$$\begin{aligned} 8\delta \;&\geq\; \Big|d_2\Big(q_2,\gamma_2(a^*)\Big) \;-\; \Big|B_2(q_2) \;-\; B_2\Big(\gamma_2(a^*)\Big)\Big|\Big| \\ &=\; \Big|d_2\Big(q_2,\gamma_2(a^*)\Big) \;-\; \Big|B_1(q_1) \;-\; B_1\Big(\gamma_1(a^*)\Big)\Big|\Big|. \end{aligned}$$

Since $d_1(q_1,\gamma_1(a^*)) \geq |B_1(q_1) - B_1(\gamma_1(a))|$ we obtain

$$f(a^*) \;=\; d_1\Big(q_1,\gamma_1(a^*)\Big) \;\geq\; d_2\Big(q_2,\gamma_2(a^*)\Big) \;-\; 8\delta \;\geq\; g(a^*) \;-\; 8\delta.$$

Since $g|_{[0,a^*]}$ and $g|_{[a^*,a^*+b^*]}$ are 12δ-T-functions and f is a δ'-T-function we see that $(g-f)_+ \leq 20\delta + \delta'$ which implies that $\max\{f,g\}$ is a $20\delta + 2\delta'$-T-function.
$\square$



Figure 4: This figure visualizes the situation in the proof of Proposition 3.

We finally indicate how the arguments of this section have to be modified to prove Theorem 1. In the case of Theorem 1 let $B_1 := d(z_1,\cdot)$, $B_2 := d(z_2,\cdot)$ and

$z = (z_1, z_2)$. The B_i-rays correspond to the geodesics $\gamma_i = \gamma_{p_i z_i}$, $\gamma'_i = \gamma_{p'_i z_i}$. Let further

$$\begin{array}{rcll} \gamma : [0, d(p,z)] & \longrightarrow & Y\ , & \qquad \gamma = (\gamma_1, \gamma_2) \\ \gamma' : [0, d(p,z)] & \longrightarrow & Y\ , & \qquad \gamma' = (\gamma'_1, \gamma'_2)\ , \end{array}$$

$a_i := (p'_i \cdot z_i)_{p_i}$ and $b_i := (p_i \cdot z_i)_{p'_i}$.
While the definition of Γ is analog to the one in the case of Theorem 2, the definition of $\Gamma^c_{pp'}$ needs to be slightly modified in the case that $a + 2\delta > d(p,z)$. In that case just take

$$\Gamma^c_{pp'} \ :=\ \gamma|_{[0,a+\tau]} \ *\ \gamma|_{[0,b+\tau]}{}^{-1},$$

where $a \leq \tau < 2\delta$ is chosen such that $\gamma(\tau) = \gamma'(\tau) = z$.
The proof of Lemma 8 stays valid in the case $y_0 \neq z$, $\epsilon < d(y_0, z)$. In the case $y_0 = z$ the result is obvious.

4 The boundary of Y

In the case of Theorem 1 it is easy to see that $\partial Y = \partial X_1 \times \partial X_2$. The situation of Theorem 2 is more interesting:
We study ∂Y and show that it is homeomorphic to $\partial X_1 \wedge \partial X_2$.
Recall that the Busemann functions B_i are defined as $B_i(x) = \lim_{t\to\infty}(d_i(x, \gamma_i(t)) - t)$, where $\gamma_i : [0,\infty) \longrightarrow X_i$ is a geodesic ray, $i = 1, 2$.
Let $z_i = \gamma_i(0)$ and $u_i = [\{\gamma_i(k)\}] \in \partial X_i$, $i = 1, 2$. Let further $z = (z_1, z_2) \in Y$ and $\gamma(t) = (\gamma_1(t), \gamma_2(t))$. Then γ is a ray with $\gamma(0) = z$ and clearly the Busemann function $B : Y \longrightarrow \mathbb{R}$ of this ray is $B(y_1, y_2) = B_1(y_1) = B_2(y_2)$.
Let $u := [\{\gamma(k)\}] \in \partial Y$. Using the results of Sections 2 and 3 there exists a Δ such that the following holds:

(1) Y is Δ-hyperbolic,

(2) $d(\Gamma^*_{xy}(t), \gamma_{xy}(t)) \leq \Delta$,

(3) $d(\Gamma^*_{xy}(t), \Gamma^*_{xz}(t)) \leq \Delta \quad \forall\, 0 \leq t \leq (y \cdot z)_x$,

(4) $d(\Gamma^*_{zx}(t), \gamma(t)) \leq \Delta \quad \forall\, 0 \leq t \leq (x \cdot B)_z$,

(5) $|d_m(x,y) - d(x,y)| \leq \Delta$.

For any point $v \in \partial Y$ consider a geodesic ray $\sigma : [0,\infty) \longrightarrow Y$ such that $\sigma(0) = z$ and $v = [\{\sigma(k)\}]$. Consider the curves $\Gamma^*_{z\sigma(k)}$ with $d(\Gamma^*_{x\sigma(k)}(t), \sigma(t)) \leq \Delta$ for $0 \leq t \leq k$. By Section 3 $\Gamma^*_{z\sigma(k)} = \gamma|_{[0,a^*_k]} * \gamma^k|_{[0,b^*_k]}{}^{-1}$, where $|a^*_k - (\sigma \cdot B)_z| \leq \Delta$ and $\gamma^k_i : [0,\infty) \longrightarrow X_i$ is a B_i-ray with $\gamma^k_i(0) = \sigma_i(k)$ and $d(\gamma^k_i(b^*_k), \gamma_i(a^*_k)) \leq \Delta$.
We distinguish two cases:
(Case 1) There exists a subsequence $\{a^*_{k_j}\}$ *with* $\lim_{j\to\infty}$.
Then $\lim_{j\to\infty}(\sigma(k_j) \cdot B)_z = \infty$ and thus $\liminf_{j,l\to\infty}(\sigma(k_j) \cdot \gamma(l))_z = \infty$, which

implies $\{\sigma(k_j)\} \sim \{\gamma(j)\}$, hence $[\{\sigma(k_j)\}] = u$. Since $\sim$ is an equivalence relation and clearly $\{\sigma(k_j)\} \sim \{\sigma(j)\}$ we also see that $\lim_{k\to\infty} a_k = \infty$.
(Case 2) $\{a_k^*\}$ *is bounded.*
Then it holds $b_k^* \longrightarrow \infty$. Reparameterize $\gamma_i^k : [0,\infty) \longrightarrow X_i$ as

$$\bar{\gamma}_i^k : [a_k^* - b_k^*, \infty)\ , \qquad \bar{\gamma}_i^k(t) \ = \ \gamma_i^k(t + b_k^* - a_k^*).$$

By the discussion of Sections 2 and 3 the $\bar{\gamma}_i^k$ converge to a complete geodesic γ_i^* with $B_i(\gamma_i(t)) = -t$ and $d_i(\gamma_i(a_k^*), \gamma^*(a_k^*)) \leq 8\delta$. Clearly we have $[\{\gamma_i^*(k)\}] = u_i$. Let $v_i := [\{\gamma_i^*(-k)\}] \in \partial X_i \setminus \{u_i\}$.

From the discussion of Case 2 it is not difficult to show that the map

$$\begin{array}{ccc} \partial Y \setminus \{u\} & \longrightarrow & (\partial X_1 \setminus \{u_1\}) \ \times \ (\partial X_2 \setminus \{u_2\}) \\ & & \\ v & \longmapsto & (v_1, v_2) \end{array}$$

is a homeomorphism, which by the discussion of Case 1 extends naturally to a homeomorphism

$$\partial Y \ \longrightarrow \ \partial X_1 \ \wedge \ \partial X_2 .$$

UNIVERSITÄT ZÜRICH, MATHEMATISCHES INSTITUT, WINTERTHURERSTRASSE 190, CH-8057 ZÜRICH, SWITZERLAND
E-mail addresses: foertsch@math.unizh.ch vschroed@math.unizh.ch

References

[BeKa] N.Benakli & I. Kapovich, *Boundaries of hyperbolic groups*, preprint math.GR/0202286

[BrFa] N. Brady & B. Farb, *Filling-Invariants at Infinity for Manifolds of Nonpositive Curvature*, Trans. Amer. Math. Soc., Vol. 350, Num. 8, 1998

[BriH] M. Bridson & A. Haefliger, *Metric spaces of non-positive curvature*, Springer Verlag Berlin 1999

[BuBuI] D. Burago & Y. Burago & S. Ivanov, *A course in Metric Geometry*, Graduate Studies in Mathematics, Vol. 33, Amer. Math. Soc., 415pp, 2001

[F1] T. Foertsch, *Bilipschitz Embeddings of Negative Sectional Curvature in Products of Warped Product Manifolds*, Proc. Amer. Math. Soc. 130, 2089-2096, (2002)

[F2] *The Hyperbolic Rank of Homogeneous Hadamard Manifolds*, to appear in the Manuscripta Mathematicae

[FS] T. Foertsch & V. Schroeder, *Hyperbolic Rank of Products*, preprint

[L1] E. Leuzinger, *Corank and Asymptotic Filling-Invariants for Symmetric Spaces*, Geom. Funct. Anal., vol.10, no4, 863-873, (2000)

[L2] *Bi-Lipschitz Embeddings of Trees into Euclidean Buildings*, preprint

[M] C.R.F. Maunder *Algebraic Topology*, Cambridge University Press, 1980